

# Exact Order of Subsets of Asymptotic Bases in Additive Number Theory

XING-DE JIA

*Department of Mathematics, Graduate School and University Center,  
City University of New York, New York, New York 10036*

*Communicated by P. Erdős*

Received April 23, 1987

Let  $A$  be an asymptotic basis of order  $h$ . Define

$$I_k(A) = \{F \mid F \subseteq A, |F| = k \text{ and } A \setminus F \text{ is a basis}\},$$

where  $|F|$  indicates the number of elements in the finite set  $F$ . We denote by  $g(A)$  the exact order of  $A$ . Define

$$G_k(h) = \max_{\substack{A \\ g(A) \leq h}} \max_{F \in I_k(A)} g(A \setminus F).$$

An open problem in additive number theory is to calculate  $G_k(h)$ . We prove in this paper that

$$G_k(h) \geq (4/3)(h/(k+1))^{k+1} + O(h^k)$$

as  $h$  tends to infinity, which is an improvement of a result of Melvyn B. Nathanson (The exact order of subsets of additive bases, in "Proceedings, New York Number Theory Seminar, 1982," Lecture Notes in Mathematics, Vol. 1052, pp. 273–277, Springer-Verlag, 1984). On the other hand, we estimate  $G_k(h)$  as  $k$  tends to infinity for any fixed integer  $h$ . It is proved that  $G_k(h)$  has order of magnitude  $k^{h-1}$  as  $k$  tends to infinity for any fixed  $h \geq 2$ . © 1988 Academic Press, Inc.

## 1. INTRODUCTION

Let  $A$  be a set of integers, and let  $hA$  denote the set of all sums of  $h$  not necessarily distinct elements of  $A$ . If  $hA$  contains all sufficiently large integers, then  $A$  is an *asymptotic basis* of order  $h$ . The set  $A$  is an asymptotic basis if  $A$  is an asymptotic basis of order  $h$  for some  $h$ . If  $A$  is an asymptotic basis, the *exact order* of  $A$ , denoted  $g(A)$ , is the smallest integer  $h$  such that  $A$  is an asymptotic basis of order  $h$ .

Let  $A$  be an asymptotic basis,  $a \in A$ . It is not necessary that  $A \setminus \{a\}$  is also an asymptotic basis. We denote by  $I$  the set of elements  $a \in A$  such that

$A \setminus \{a\}$  is an asymptotic basis. For any  $a \in I$ , Erdős and Graham [1] investigated how large  $g(A \setminus \{a\})$  can be in terms of  $g(A)$ . Let

$$G_1(h) = \max_{\substack{A \\ g(A) \leq h}} \lim_{a \in I} g(A \setminus \{a\}).$$

Erdős and Graham [1] proved in 1980 that

$$\frac{1}{4}(1 + o(1)) h^2 \leq G_1(h) \leq (5/4)(1 + o(1)) h^2.$$

In his doctoral thesis, G. Grekos [2] improved this estimate to

$$\frac{1}{3}h^2 + O(h) \leq G_1(h) \leq h^2 + h,$$

where the upper bound holds for any  $h > 1$ . Recently, Nash obtained in his doctoral thesis [6] that

$$G_1(h) \leq \frac{1}{2}h^2 + O(h).$$

However, it is an open problem to improve the estimates of  $G_1(h)$  (see [7] or [6]).

In 1982, Nathanson considered the general form of this problem. Instead of cancelling only one element in  $A$ , he cancelled a finite subset of  $A$  with fixed cardinality.

Let  $k \geq 1$  be an integer. If  $A$  is an asymptotic basis, let  $I_k(A)$  denote the set of all subsets  $F \subseteq A$  such that  $F$  has cardinality  $k$  and the set  $A \setminus F$  is an asymptotic basis. Define

$$G_k(h) = \max_{\substack{A \\ g(A) \leq h}} \max_{F \in I_k(A)} g(A \setminus F).$$

Nathanson [7] proved that

$$G_k(h) \geq ([h/(k+1)] + 1)^{k+1} - 1,$$

where  $h > k$  and  $[x]$  indicates the largest integer not exceeding  $x$ . In Section 2, we improve this estimate to

$$G_k(h) \geq \frac{4}{3}(h/(k+1))^{k+1} + O(h^k).$$

In case  $k = 1$ , this is the lower bound of Grekos.

Professor Nathanson [7] thinks that it is significant to estimate  $G_k(h)$  as  $k$  tends to infinity for any fixed integer  $h \geq 2$ . Nash [5] proved that  $G_k(2) = 2k + 2$  for any  $k \geq 1$ . In Section 3, we prove that  $G_k(h)$  has order of magnitude  $k^{h-1}$  as  $k$  tends to infinity for any fixed  $h \geq 2$ , namely

$$G_k(h) + 1 \geq \frac{2}{(h-1)^{h-1}} k^{h-1} + \frac{4h-5}{(h-1)^{h-2}} k^{h-2} + O(k^{h-3}),$$

$$G_k(h) + 1 \leq \frac{2}{(h-1)!} k^{h-1} + \frac{h-1}{(h-2)!} k^{h-2} + O(h^{h-3}).$$

If  $h = 2$  this is the result of Nash.

## 2. ESTIMATE OF THE LOWER BOUND OF $G_k(h)$ FOR ANY $k$

LEMMA 2.1. *Let  $h \geq k + 1$  and  $k \geq 1$ , and*

$$u = [h/(k+1)], \quad u' = [2h/(3k+3)],$$

$$\sigma = h - (k-1)u - 2u',$$

$$b_1 = [2h/(k+1)],$$

$$b_i = ub_{i-1} + \sigma, \quad \text{for } i = 1, 2, \dots, k,$$

$$d = u'b_k + b_1 - h + (k-1)u + u'.$$

Then

$$(i) \quad md + \sum_{i=1}^k x_i b_i + h - \sum_{i=1}^k x_i \geq (m-1)d + \sum_{i=1}^k x_i b_i + b_1 + u'b_k;$$

$$(ii) \quad (m-1)d + \sum_{i=1}^k x_i b_i + u'b_k + h - \sum_{i=1}^k x_i - u' \geq md + \sum_{i=1}^k x_i b_i$$

hold for any  $m \geq 1$ , and

$$0 \leq x_i \leq u \quad \text{for } i = 1, 2, \dots, k-1, \quad 0 \leq x_k \leq u'.$$

*Proof.* From the definitions, we have

$$\begin{aligned} & md + \sum_{i=1}^k x_i b_i + h - \sum_{i=1}^k x_i \\ &= (m-1)d + \sum_{i=1}^k x_i b_i + h - \sum_{i=1}^k x_i + d \\ &= (m-1)d + \sum_{i=1}^k x_i b_i + b_1 + u'b_k + \sum_{i=1}^{k-1} (u - x_i) + (u' - x_k) \\ &\geq (m-1)d + \sum_{i=1}^k x_i b_i + b_1 + u'b_k, \end{aligned}$$

which shows (i). It follows from the definitions that

$$\begin{aligned}
 & (m-1)d + \sum_{i=1}^k x_i b_i + u' b_k + h - \sum_{i=1}^k x_i - u' \\
 &= md - u' b_k - b_1 + h - (k-1)u - u' + \sum_{i=1}^k x_i b_i \\
 &\quad + u' b_k + h - \sum_{i=1}^k x_i - u' \\
 &= md + \sum_{i=1}^k x_i b_i + 2h - (k-1)u - 2u' - \sum_{i=1}^k x_i - b_1 \\
 &\geq md + \sum_{i=1}^k x_i b_i + 2h - 2(k-1)u - 3u' - b_1.
 \end{aligned}$$

Observing that

$$\begin{aligned}
 2(k-1)u + 3u' + b_1 &\leq 2(k-1)\frac{h}{k+1} + 3\frac{2h}{3k+3} + \frac{2h}{k+1} \\
 &= 2h\left(\frac{k-1}{k+1} + \frac{1}{k+1} + \frac{1}{k+1}\right) \\
 &= 2h,
 \end{aligned}$$

we have

$$(m-1)d + \sum_{i=1}^k x_i b_i + 2h - 2(k-1)u - 3u' - b_1 \geq md + \sum_{i=1}^k x_i b_i,$$

which implies (ii). The proof is complete.

**LEMMA 2.2.** *Let  $h, k, u, u', b_1, \dots, b_k$  be as in Lemma 1.1. If*

$$0 \leq x_i \leq u \quad \text{for } i = 1, 2, \dots, k-1, \quad 0 \leq x_k \leq 2u',$$

then

$$\sum_{i=\lambda+1}^k x_i b_i + u b_\lambda + h - \sum_{i=\lambda+1}^k x_i - u \geq \sum_{i=\lambda+1}^k x_i b_i + b_{\lambda+1}$$

holds for any  $1 \leq \lambda \leq k-1$ .

*Proof.* Noting that  $\lambda \geq 1$ , we have

$$\begin{aligned}
 h - \sum_{i=\lambda+1}^k x_i - u &\geq h - (k-2)u - 2u' - u \\
 &= h - (k-1)u - 2u' \\
 &= \sigma.
 \end{aligned}$$

Therefore

$$\begin{aligned} \sum_{i=\lambda+1}^k x_i b_i + u b_\lambda + h - \sum_{i=\lambda+1}^k x_i - u &\geq \sum_{i=\lambda+1}^k x_i b_i + u b_\lambda + \sigma \\ &= \sum_{i=\lambda+1}^k x_i b_i + b_{\lambda+1}, \end{aligned}$$

which proves the lemma.

LEMMA 2.3. *Let  $d$  be as in Lemma 1.1. Then*

$$D = \{a \mid a \equiv 0 \text{ or } 1 \pmod{d}\}$$

*is an asymptotic basis with*

$$g(D) = \frac{4}{3}(h/(k+1))^{k+1} + O(h^k).$$

*Proof.*  $h \geq 3k+3$  implies  $d > 2$ . Let  $n$  be any positive integer with

$$n = qd + r,$$

where  $0 \leq r < d$ . If  $r = 0$  then  $n = qd \in D \subseteq (d-1)D$ . If  $r > 0$  then

$$n = (qd+1) + (r-1) \in rD \subseteq (d-1)D.$$

Hence  $D$  is an asymptotic basis with the exact order  $g(D) < d$ .

For any  $s$ , if

$$sd + (d-1) = \sum_{i=1}^t a_i, \quad a_i \in D,$$

then it follows from the definition of  $D$  that there exist at least  $d-1$  elements  $a_i$  of  $\{a_1, a_2, \dots, a_t\}$  such that  $a_i \equiv 1 \pmod{d}$ , which implies that  $t \geq d-1$ . Therefore

$$\begin{aligned} g(D) &= d-1 \\ &= u'b_k + b_1 - h + (k-1)u + u' - 1 \\ &= u'ub_{k-1} + u'\sigma + b_1 - h + (k-1)u + u' - 1 \\ &= \dots \\ &= u'u^{k-1}b_1 + u'(u^{k-2} + \dots + u + 1) + b_1 - h + (k-1)u + u' - 1 \\ &= \frac{2h}{3k+3} \left( \frac{h}{k+1} \right)^{k-1} \frac{2h}{k+1} + O(h^k) \\ &= \frac{4}{3} \left( \frac{h}{k+1} \right)^{k+1} + O(h^k). \end{aligned}$$

The proof is complete.

THEOREM 2.4. *Let  $k \geq 1$ . Then*

$$G_k(h) \geq \frac{4}{3} \left( \frac{h}{k+1} \right)^{k+1} + O(h^k)$$

as  $h$  tends to infinity.

*Proof.* Taking

$$F = \{b_1, b_2, \dots, b_k\}, \quad A = F \cup D,$$

where  $b_1, b_2, \dots, b_k, d$ , and  $D$  are as in Lemmas 1.1 and 1.2, we obtain with Lemma 1.3 that  $A$  is an asymptotic basis. Now we assume  $h \geq 3k + 3$ . Then

$$4 \leq b_1 < b_2 < \dots < b_k < d,$$

which shows  $A \setminus F = D$ , so that  $F \in I_k(A)$ . By Lemma 1.3, it is sufficient to prove  $g(A) \leq h$ .

Let  $m \geq$  and

$$0 \leq x_i < u \quad \text{for } i = 1, 2, \dots, k-1, \quad 0 \leq x_k \leq u'.$$

Since

$$\begin{aligned} 2 + \sum_{i=1}^k x_i + u' &\leq 2 + (k-1)u + 2u' \\ &\leq 2 + (k-1)h/(k+1) + 4h/(3k+3) \\ &= h - (2h/(3k+3) - 2) \\ &\leq h, \end{aligned}$$

we have

$$\begin{aligned} md + \sum_{i=1}^k x_i b_i &\in hA, \\ (m-1)d + \sum_{i=1}^k x_i b_i + b_1 + u'b_k &\in hA. \end{aligned}$$

Thus, any  $x$  satisfying

$$md + \sum_{i=1}^k x_i b_i \leq x \leq md + \sum_{i=1}^k x_i b_i + h - \sum_{i=1}^k x_i$$

or

$$\begin{aligned} (m-1)d + \sum_{i=1}^k x_i b_i + b_1 + u'b_k &\leq x \\ &\leq (m-1)d + \sum_{i=1}^k x_i b_i + b_1 + u'b_k + h - \sum_{i=1}^k x_i - 1 - u' \end{aligned}$$

is contained in  $hA$ . It therefore follows from Lemma 1.1 that any  $x$  satisfying that

$$md + \sum_{i=1}^k x_i b_i \leq x \leq (m-1)d + \sum_{i=1}^k x_i b_i + b_1 + u'b_k - \sum_{i=1}^k x_i - 1 - u'$$

is contained in  $hA$ . But

$$(m-1)d + \sum_{i=1}^k x_i b_i + b_1 + u'b_k + h - \sum_{i=1}^k x_i - 1 - u' \geq md + \sum_{i=1}^k x_i b_i + b_1,$$

we therefore obtain that any  $x$  satisfying

$$md + \sum_{i=1}^k x_i b_i \leq x \leq md + (x_1 + 1)b_1 + \sum_{i=2}^k x_i b_i$$

is already contained in  $hA$ . The arbitrariness of  $x_1$  implies that  $hA$  contains all integers in the interval

$$\left[ md + \sum_{i=2}^k x_i b_i, md + ub_1 + \sum_{i=2}^k x_i b_i \right].$$

Using Lemma 1.2, we have

$$md + \sum_{i=2}^k x_i b_i + ub_i + h - \sum_{i=2}^k x_i - u \geq md + (x_2 + 1)b_2 + \sum_{i=3}^k x_i b_i,$$

which implies that any  $x$  satisfying that

$$md + \sum_{i=2}^k x_i b_i \leq x \leq md + (x_2 + 1)b_2 + \sum_{i=3}^k x_i b_i$$

is contained in  $hA$ . Again by the arbitrariness of  $x_2$ , we have that  $hA$  contains all integers in the interval

$$\left[ md + \sum_{i=3}^k x_i b_i, md + ub_2 + \sum_{i=3}^k x_i b_i \right].$$

Again using Lemma 1.2, we have that  $hA$  contains all integers in the interval

$$\left[ md + \sum_{i=3}^k x_i b_i, md + (x_3 + 1)b_3 + \sum_{i=4}^k x_i b_i \right].$$

Using a similar argument, we obtain that any  $x$  satisfying

$$md \leq x \leq md + u'b_k$$

is contained in  $hA$ . Observing

$$\begin{aligned} b_1 - h + (k-1)u + u' &\leq \frac{2h}{k+1} - h + (k-1)\frac{h}{k+1} + \frac{2h}{3k+3} \\ &\leq \frac{2h}{3k+3} < h - u' - 1, \end{aligned}$$

we finally obtain that any  $x$  satisfying  $md \leq x \leq (m+1)d$  is contained in  $hA$  for any  $m > 1$ . This means that  $x \in hA$  for all  $x > d$ , i.e.,  $g(A) \leq h$ , which ends the proof.

### 3. ESTIMATE OF $G_k(h)$ FOR GIVEN $h$

In this section,  $[a, b]$  denotes the set of all integers  $n$  such that  $a \leq n \leq b$ .

LEMMA 3.1. Let  $h \geq 2$ ,  $k \geq h-1$ . Let  $l = [k/(h-1)]$ . Define

$$\begin{aligned} b_0 &= 1, \\ b_{il+j} &= b_{il} + j \left( 2 + \sum_{r=1}^i b_{rl} \right), \end{aligned}$$

for  $i = 0, 1, \dots, h-2$  and  $j = 1, 2, \dots, l$ . Let  $m \geq 2$ . Let

$$D = \{0, 1, md, md+1\}.$$

Then

$$\left[ md + 2 + \sum_{r=1}^{i-1} b_{rl}, md + 1 + \sum_{r=1}^i b_{rl} \right] \subseteq (i+1)(\{b_1, \dots, b_{il}\} \cup D) \quad (1)$$

holds for  $i = 1, 2, \dots, h-1$ .

*Proof.* If  $n \in [md+2, md+1+b_1]$ , then

$$2 \leq n - md \leq b_1 + 1.$$

If  $n - md$  is odd, then there exists a  $b_j$  ( $1 \leq j \leq l$ ) such that

$$b_j = n - md,$$

i.e.,  $a = md + b_j$ . If  $n - md$  is even, then  $n = (md+1) + 1$  or there exists a  $b_j$  such that  $n = (md+1) + b_j$ . Hence

$$n \in 2(\{b_1, \dots, b_1\} \cup D).$$



Now assume (1) holds for any  $i \leq s-1$ , it follows that for any  $i \leq s-1$ ,

$$\left[ md, md+1 + \sum_{r=1}^i b_{r_l} \right] \subseteq (i+1)(\{b_1, \dots, b_{il}\} \cup D).$$

Let

$$n \in \left[ md+2 + \sum_{r=1}^{s-2} b_{r_l}, md+1 + \sum_{r=1}^s b_{r_l} \right].$$

If

$$md+2 + \sum_{r=1}^{s-1} b_{r_l} \leq n \leq md + b_{(s-1)l} + 1 - 1,$$

then

$$md+2 + \sum_{r=1}^{s-2} b_{r_l} \leq n - b_{(s-1)l} \leq md+1 + \sum_{r=1}^{s-1} b_{r_l}.$$

It follows from the assumption that

$$n - b_{(s-1)l} \in (\{b_1, \dots, b_{(s-1)l}\} \cup D),$$

which means  $n \in (s+1)(\{b_1, \dots, b_{(s-1)l}\} \cup D)$ .

If there exists a  $j: 1 \leq j \leq l-1$  such that

$$md + b_{(s-1)l+j} \leq n \leq md + d_{(s-1)l+j} + 1 - 1,$$

then

$$md \leq n - b_{(s-1)l+j} \leq md+1 + \sum_{r=1}^{s-1} b_{r_l}.$$

It follows from the assumption that

$$n \in (s+1)(\{b_1, \dots, b_{(s-1)l}\} \cup D).$$

If

$$md + b_{sl} \leq n \leq md+1 + \sum_{r=1}^s b_{r_l},$$

then  $n - b_{sl} \in s(\{b_1, \dots, b_{(s-1)l}\} \cup D)$ , thus

$$n \in (s+1)(\{b_1, \dots, b_{(s-1)l}\} \cup D).$$

The proof is complete.

The notation  $hA$  can be generalized. Let  $A_1, \dots, A_r$  be sets of integers. Define

$$\sum_{i=1}^r A_i = \left\{ \sum_{i=1}^r a_i \mid a_i \in A_i \text{ for } i = 1, 2, \dots, r \right\}.$$

Let  $B$  be a set of integers, and let  $g$  be a positive integer.  $B^{(g)}$  denotes the set of nonnegative integers congruent to some element of  $B$ . The notation  $A \sim B$  means

$$A \cap \{n \geq N \mid n \in \mathbb{N}\} = B \cap \{n \geq N \mid n \in \mathbb{N}\}$$

for some  $N$ .  $A(m)$  is the size of the set  $\{a \in A \mid a \leq m\}$ . The lower density of  $A$  is defined as

$$dA = \liminf_{m \rightarrow \infty} \frac{A(m)}{m}.$$

To give an upper bound for  $G_k(h)$ , we need the following Kneser's Theorem.

**KNESER'S THEOREM** (see [4]). *Let*

$$C = \sum_{i=1}^n A_i.$$

*Then either*

$$dC \geq \sum_{i=1}^n dA_i$$

*or*

$$C \sim C^{(g)}$$

*for some  $g$ .*

Now we give an upper bound for  $G_k(h)$ .

**LEMMA 3.2.** *Let  $h \geq 2$ . Then*

$$G_k(h) + 1 \leq 2k^{h-1}/(h-1) + (h+1)k^{(h-2)/(h-2)} + O(k^{h-3})$$

*as  $k$  tends to infinity.*

*Proof.* Let  $A$  be an asymptotic basis of order  $h$ . Let  $F \in I_k(A)$ . Let  $B = A \setminus F$ . We have

$$hB \cup F + (h-1)B \cup \dots \cup (h-1)F + B \cup hF \sim N. \quad (2)$$

Hence  $dhB > 0$  and

$$dhB + kd(h-1)B + \binom{k+1}{2}d(h-2)B + \cdots + \binom{k+h-2}{h-1}dB \geq 1.$$

Let  $k \geq \max(h-1, 2)$ . Then we have

$$\binom{k+h-1}{k} - 1 \geq h.$$

Therefore

$$\begin{aligned} & d\left(\binom{k+h-1}{k} + 1\right)B + dhB + kd(h-1)B \\ & + \binom{k+1}{2}d(h-2)B + \cdots + \binom{k+h-2}{h-1}dB > 1. \end{aligned}$$

It follows from Kneser's theorem that we have a  $g$  such that

$$\begin{aligned} & \left(\binom{k+h-1}{k} - 1 + \sum_{i=0}^{h-1} \binom{k+i-1}{i} (h-i)\right)B \\ & \sim \left(\binom{k+h-1}{k} - 1 + \sum_{i=0}^{h-1} \binom{k+i-1}{i} (h-i)\right)B^{(g)}. \end{aligned}$$

Take the smallest  $g$  for which this last relation holds. We show that  $g = 1$ .

It follows from (2) that

$$hB^{(g)}(g) + k((h-1)B^{(g)}(g)) + \cdots + \binom{k+h-2}{h-1}(B^{(g)}(g)) \geq g.$$

Using Kneser's theorem, we have

$$\left(\sum_{i=0}^{h-1} \binom{k+i-1}{i} (h-i)\right)B^{(g)}(g) \geq g - \binom{h+k-1}{k} + 1.$$

It is clear that if  $nB^{(g)}(g) < g$ , then  $(n+1)B^{(g)}(g) > nB^{(g)}(g)$ . Therefore

$$\left(\binom{h+k-1}{k} - 1 + \sum_{i=0}^{h-1} \binom{k+i-1}{i} (h-i)\right)B^{(g)}(g) = g.$$

This implies  $g = 1$ . Therefore

$$\begin{aligned} G_k(h) + 1 & \leq \binom{h+k-1}{k} + \sum_{i=0}^{h-1} \binom{k+i-1}{i} (h-i) \\ & = \frac{2}{(h-1)!} k^{h-1} + \frac{h+1}{(h-2)!} k^{h-2} + O(k^{h-3}) \end{aligned}$$

as  $k$  tends to infinity. This completes the proof.

THEOREM 3.3. *Let  $h \geq 2$ . Then*

$$\begin{aligned} G_k(h) + 1 &\geq 2(k/(h-1))^{h-1} + (4h-5)(k/(h-1))^{h-2} + O(k^{h-3}); \\ G_k(h) + 1 &\leq 2k^{h-1}/(h-1)! + (h-1)k^{h-2}/(h-2)! + O(k^{h-3}) \end{aligned} \quad (3)$$

as  $k$  tends to infinity. In particular,  $G_k(2) = 2k + 2$  holds for any  $k \geq 1$ .

*Proof.* The second inequality has already been established in Lemma 3.2. We now prove the first one.

Suppose  $k \geq h-1$ . Let  $1, b_0, \dots, b_{(h-1)l}$  be as in Lemma 3.1. Define

$$b_{(h-1)l+j} = b_{(h-1)l} + j \left( 2 + \sum_{r=1}^{h-2} b_{rl} \right)$$

for  $j = 1, 2, \dots, k - (h-1)l$ . Define

$$d = b_k + 2 + \sum_{r=1}^{h-2} b_{rl}.$$

Let

$$\begin{aligned} F &= \{b_1, b_2, \dots, b_k\}, \\ A &= F \cup \{a \in \mathbb{N} \mid a \equiv 0 \text{ or } 1 \pmod{d}\}. \end{aligned}$$

It is clear that both  $A$  and  $A \setminus F$  are asymptotic bases.

We prove that  $A$  is of order  $h$ . Let  $m \geq 0$  be any integer. By Lemma 3.1 we have

$$\left[ md + 2 + \sum_{r=1}^{s-1} b_{rl}, md + 1 + \sum_{r=1}^s b_{rl} \right] \subseteq (s+1)A$$

for  $s = 1, 2, \dots, h-1$ , which implies that

$$\left[ md, md + 1 + \sum_{r=1}^{h-1} b_{rl} \right] \subseteq hA.$$

Let

$$n \in \left[ md + 2 + \sum_{r=1}^{h-1} b_{rl}, md + d - 1 \right].$$

Suppose

$$md + b_{(h-1)l} + t \left( 2 + \sum_{r=1}^{h-2} b_{rl} \right) \leq n < md + b_{(h-1)l} + (t+1) \left( 2 + \sum_{r=1}^{h-2} b_{rl} \right)$$

for some  $t$ . Then

$$n - b_{(h-1)l+1} = n - \left( (b_{(h-1)l} + t \left( 2 + \sum_{r=1}^{h-2} b_{rl} \right) \right)$$

is contained in  $[md, md + 1 + \sum_{r=1}^{h-2} b_{rl}]$ , hence in  $(h-1)A$ . This means  $n \in hA$ . Therefore  $[md, (m+1)d-1] \subseteq hA$ . Consequently,  $g(A) \leq h$ .

Since  $A \setminus F = \{a \in N \mid a \equiv 0 \text{ or } 1 \pmod{d}\}$ , it follows that  $g(A \setminus F) = d-1$ . Hence  $G_k(h) + 1 \geq d$ . From the definition, a simple calculation shows that

$$b_{sl} = lb_{(s-1)l} + (b_{(s-1)l} + lb_{(s-2)l}) + O(l^{s-2})$$

holds for  $s = 2, 3, \dots, h-1$ . This implies that

$$b_{(h-1)l} = 2l^{h-1} + (4h-7)l^{h-2} + O(l^{h-3}).$$

Therefore

$$\begin{aligned} G_k(h) + 1 &\geq d = b_k + 2 + \sum_{r=1}^{h-2} b_{rl} \\ &= b_{(h-1)l} + (k - (h-1)l) \left( 2 + \sum_{r=1}^{h-2} b_{rl} \right) + 2 + \sum_{r=1}^{h-2} b_{rl} \\ &= 2l^{h-1} + (4h-7)l^{h-2} + b_{(h-2)l} + 2 + \sum_{r=1}^{h-3} b_{rl} \\ &\quad + (k - (h-1)l) \left( 2 + \sum_{r=1}^{h-2} b_{rl} \right) + O(l^{h-3}) \\ &\geq 2l^{h-1} + (4h-5)l^{h-2} + 2l^{h-2}(k - (h-1)l) + O(l^{h-3}) \\ &\geq 2(k/(h-1))^{h-1} + (4h-5)(k/(h-1))^{h-2} \\ &\quad - 2(h-1)(k/(h-1)-1)(k/(h-1))^{h-2} \\ &\quad + 2(k - (h-1)l)(k/(h-1))^{h-2} + O(k^{h-3}) \\ &= 2(k/(h-1))^{h-1} + (4h-5)(k/(h-1))^{h-2} + O(k^{h-3}), \end{aligned}$$

which proves the inequality.

From Lemma 3.2 and the argument above, we see that the lower bound and the upper bound are polynomials of  $k$  with integral coefficients. Hence when  $h=2$ , the  $O$ 's in (3) are zero, namely,  $G_k(2) = 2k + 2$ . It is readily seen that this holds for any  $k \geq 2$ , hence for any  $k \geq 1$  since  $G_1(2) = 4$ . (see [1]). The proof is complete.

## ACKNOWLEDGMENTS

I thank Professor M. B. Nathanson and Professor Pin-tsung Shao for their helpful suggestions and Dr. J. C. M. Nash for sending me a copy of his doctoral thesis.

## REFERENCES

1. ERDÖS AND R. L. GRAHAM, On bases with an exact order, *Acta Arith.* **37** (1980), 201–207.
2. G. GREKOS, “Quelques Aspects de la Théorie Additive des Nombres,” Thesis, Université de Bordeaux I, 1982.
3. H. HALBERSTAM AND K. F. ROTH, “Sequences,” Oxford Univ. Press, Oxford, 1966.
4. H. B. MANN, “Addition Theorems,” Interscience Pub., New York, 1965.
5. J. C. M. NASH, “Results in Bases in Additive Number Theory,” Thesis, Rutgers University, New Jersey, 1985.
6. J. C. M. NASH AND M. B. NATHANSON, Cofinite subsets of asymptotic bases for the positive integers, *J. Number Theory*, **20** (1985), 363–372.
7. M. B. NATHANSON, The exact order of subsets of additive bases, in “Proceedings, Number Theory Seminar, 1982,” Lecture Notes in Mathematics, Vol. 1052, pp. 273–277, Springer-Verlag, 1984.